

Preparation for Bowel Surgery: MoviPrep

Date_	/	/	_ Day of Surge	ery: Arrive at:	am/pm
Pleas	se check in	at PACE	E/Day Stay located at 8	00 Prudential D	rive, 1 st Floor of the
	<u>Hea</u>	rt Hospit	al_		
Pleas • • • •	Drink 3 pro throughou Your eveni Do NOT di Do NOT sr	balanced of the control of the day of the day of the day of the control of the co	ow: diet starting 6 weeks pri es (Ensure, Enlive, Ens prior to surgery should be CLEAR LIQUI ol for at least 24 hours b e/chew tobacco at least op at least 6 weeks prior	sure Plus or Boos DS ONLY before surgery 24 hours before	
Your	prescriptio	ns have l	been e-prescribed to:		
Pleas	e follow the	CHECKE	D prep below		
water Ensui	liquids are: \and clear b	Water, Je roth. You	liquids at 8am the day place, grape juice, apple justing should drink 3 protein son Boost Plus. Abstair	uice, popsicles, t hakes throughou	olack coffee, tea, at the day such as
	1pm Take	the first o	dose of antibiotics		
	2pm Take	the seco	nd dose of antibiotics		
	MoviPrep	starting a	t 3pm the day before yo	our surgery	

- Empty one "pouch A" and one "pouch B" into the disposable container
- Add lukewarm drinking water to the top of the container and fill to the top line
- Shake until the mix is dissolved. Do NOT add ice or any flavoring to the solution
- Refrigerate the solution
- Drink one 8 ounce glass every 10-15 minutes until the entire liter is finished
- Drink at least 16 ounces of clear liquid of your choice



- Do NOT add ice or any favoring to the solution
- Repeat the process at 6pm

DO NOT eat anything including solid foods, protein shakes, creamer, milk, gum, candy, or mints after the midnight before your procedure or surgery. You may drink only clear liquids: (water, flavored water, apple juice, cranberry juice (no pulp), black coffee (nothing added) or tea (nothing added) after midnight until 2 hours prior to your arrival time for surgery. If Baptist Health has given you a pre-surgery drink, finish contents 2 hours prior to your scheduled arrival time. Finish contents within 10 minutes. If you become nauseated, stop drinking. NOTE: Do not drink any other fluids after you finish the pre-surgery drink. If not given a pre-surgery drink, the stop drinking clear liquids 2 hours prior to scheduled arrival time. Stop drinking liquids at which is 2 hours prior to your scheduled arrival time. You can brush your teeth but don't swallow any water.

Medication to take the morning of surgery

You will be instructed at your PACE appointment which medications to take the day of surgery. Ask: 1) Which medications to take, 2) when to take them, 3) how to take them. Make sure you ask about diabetes medication, blood thinners (anticoagulants), vitamins, herbs, supplements. Depending on your situation you may need to stop taking some medications several days or weeks before surgery. If you are told to take medications the morning of surgery you may have only enough water to swallow the medication.

*** Any instructions our office provides supersedes those provided by PACE. ***

Showering Instructions

Please shower with Hibiclens (4% chlorhexidine). This skin cleanser will help reduce the concentration of bacteria on your skin before surgery and reduce your risk of infection. As indicated below, **do not** apply the Hibiclens above the level of your neck, as it can be irritating to your eyes.

Purchasing Information

Hibiclens (4% chlorhexidine) is available <u>without a prescription</u> at most retail pharmacies. Retail pharmacies may offer these products under different brand names, so be sure to verify with a pharmacist that you are selecting the correct product. **Purchase two (2) 8oz bottles.**

Instructions for Hibiclens (4% chlorhexidine shower) use:

You will perform a Hibiclens shower twice:

- 1. The night prior to your operation, and
- 2. The morning of your operation at home.



Repeat the instructions below for each Hibiclens shower.

- 1. Bathe your entire body surface (except the face and scalp). Prior to using the hibiclens you may use an antibacterial soap
- 2. Do not apply to face or scalp, as Hibiclens can irritate your eyes
- 3. You may use a clean washcloth with Hibiclens to bathe (Note: These products may permanently stain fabrics)
- 4. After rinsing thoroughly, dry off with a clean towel.
- 5. Repeat this process in the morning at home prior to coming to the hospital.
- 6. In the evening, **2 hours after your Hibiclens shower**, use the chlorhexidine wipes. *These will be provided at your PACE visit. Follow the printed instructions.*

Evening before/Morning of surgery

- Apply clean bed linens
- Wash your hair. Do not wash your hair the morning of surgery.
- Put on clean pajamas. **Do not apply any lotions or oils to the skin except for the face**
- Do NOT use a razor to shave legs or remove hair from surgical site. This could cause irritation.

After checking into the hospital, you nurse will have you apply the second application of 2% chlorhexidine wipes. The nurse will have you put on a clean hospital gown. The nurse will have you use an oral rinse to cleanse the mouth. The nurse will have you use antiseptic swabs to cleanse both nostrils ****Please let the nurse know if you have any allergies to betadine, iodine or shellfish****

Transportation

A designated driver (family member or friend) may need to drive you home upon discharge.

You cannot undergo the operation without a designated driver.

Clothing

Wear loose comfortable clothing. A minimal amount of makeup is OK. You must remove contact lenses, false eyelashes, artificial limbs, tampons, hairpins, hair decorations, rings, watches and other jewelry. You may wear dentures, eyeglasses, hearing aids and wigs to surgery, holding area.

Valuables: Please leave all valuables (rings, watches and other jewelry, etc) at home



Questions: Please call our Nurse Triage line at (904) 202-7300 Option 4 or your clinic nurse, 8 a.m. to 5 p.m. Monday through Friday and (904) 202-7300 after hours, holidays, and weekends.

Reviewed 1/2020