

Patient Education

Skin Care During and After Radiation Treatment for Breast Cancer

Radiation treatment causes side effects to your skin that are similar to those caused by sun damage. Common side effects include dryness, mild redness or tanning, then noticeable redness or tanning and then sometimes blistering.

Although every patient is different, side effects usually occur around the 2nd or 3rd week of treatment. Side effects are not usually related to your tendency to sunburn. The information below will help you care for your skin. Follow these directions, unless otherwise directed by your doctor.

General Skin Care

- Begin a skin care routine that your doctor recommends or approves.
- Aquaphor[®], Lubriderm[®], Eucerin[®] or Miaderm[®] is often recommended to patients for general skin care. Apply the product to the treatment areas **after radiation**. Avoid using products near your marking lines and stickers. It is important **not** to have thick creams on the skin at the time of treatment.
- Over the counter hydrocortisone cream can help relieve redness and itching.
- If your skin becomes tender your team may give you non-adherent medicated dressings to place over the affected areas. A member of your medical team will show you how to apply these.
- Use fragrance-free soaps with moisturizers (such as Dove[®]).
- Help prevent skin irritation by avoiding tight clothes or underwire bras, especially once you begin to have symptoms.

If these suggestions do not relieve symptoms, your medical team will discuss alternatives during your weekly doctor visit.

Treatment Marking Lines

- Treatment lines will be drawn on your skin to ensure that the radiation is aimed at the same area during each treatment. The marking ink will stain, so you may want to wear old clothing or under clothes between the marks and your good clothing. The marks may be redrawn by the therapist if they fade. When using skin care products, be careful to not rub off the lines.
- When showering, let water run over the treatment lines, avoid hot water and scrubbing.

Skin-On-Skin Contact

- Skin touching skin can happen:
 - Where our arm presses against your armpit and side of the breast.
 - Along the bottom of your breast, where it touches your abdomen.
 - Between both breasts.
- To avoid skin-on-skin contact:
 - Try to keep your arm from touching the body whenever possible.
 - Wear clothes with sleeves rather than sleeveless clothing.
 - Wear a sturdy bra without an underwire that keeps your breasts separated and lifted.
 - For large-breasted women: When you're not wearing a bra, place a soft washcloth or piece of soft cotton or flannel under your breast. Your medical team may give you non-adherent dressings to help with this.
- Dusting the breast area and inside skin folds with cornstarch helps to reduce rubbing of your skin. Apply it with a clean makeup brush or put some cornstarch into a single knee-high nylon or thin sock and knot it at the top. Gently tap the sock against the skin to dust the surface. If your doctor recommends using creams, apply those first, then dust the area with the cornstarch.

After Treatment

After your radiation treatment is complete, the skin that has been exposed to radiation may be more sensitive to the sun than before cancer. You can go out in the sun, but continue to protect your skin. Follow the guidelines below.

- Use a sunblock that is rated SPF 30 or higher on the treated area. SPF 30 absorbs about 97% of the sun's burning rays, while SPF 45 absorbs about 98%. No sunscreen or SPF will completely protect you from the sun.
- Apply the sunblock 30 minutes before you go out in the sun.
- Once you're outdoors, reapply sunscreen every two hours or more often if you're swimming or sweating.
- The treated skin will always be drier than it was before. After your symptoms have improved, moisturize everyday with the moisturizer of your choice. This will help reduce permanent tan lines.

Reviewed 10/2019